

SPEA BOMBIFRONS (Plains Spadefoot). USA: NEBRASKA: BURT Co.: NE Hwy 51, ca. 0.1 rd km E jct NE Hwy 16 (42.00275°N, 96.55379°W; WGS 84). 19 May 2017. Samuel M. Tegge and Drew R. Davis. Verified by Travis J. LaDuc. Biodiversity Collections, University of Texas at Austin (TNHC 103528 [DRD 3971]). New county record that fills a gap in the known distribution of this species in Nebraska (Ballinger et al. 2010. *Amphibians and Reptiles of Nebraska*. Rusty Lizard Press, Oro Valley, Arizona. 400 pp.; Fogell 2010. *A Field Guide to the Amphibians and Reptiles of Nebraska*. University of Nebraska–Lincoln, Lincoln, Nebraska. 158 pp.) and is known from surrounding Cuming, Dodge, Thurston, and Washington counties, Nebraska (Ballinger et al. 2010, *op. cit.*; Fogell 2010, *op. cit.*; see below) and Harrison and Monona counties, Iowa (LeClere 2013. *A Field Guide to the Amphibians and Reptiles of Iowa*. ECO Herpetological Publishing and Distribution, Rodeo, New Mexico. 349 pp.). The nearest known locality for this species is from ca. 0.5 km to the west from Cuming County (TNHC 103529; see below). Adult female (14.04 g; 50 mm SVL) collected AOR at 2157 h after heavy rains.

CUMING Co.: NE Hwy 51, ca. 0.4 rd km W jct NE Hwy 16 (42.00277°N, 96.55984°W; WGS 84). 19 May 2017. Samuel M. Tegge and Drew R. Davis. Verified by Travis J. LaDuc. TNHC 103529 (DRD 3972). New county record that fills a gap in the known distribution of this species in Nebraska (Ballinger et al. 2010, *op. cit.*; Fogell 2010, *op. cit.*) and is known from surrounding Burt, Colfax, Dodge, Stanton, Thurston, and Wayne counties, Nebraska (Ballinger et al. 2010, *op. cit.*; Fogell 2010, *op. cit.*; see above and below). The nearest known locality for this species is from ca. 0.5 km to the east in Burt County (TNHC 193528; see above). Adult female (10.25 g; 49 mm SVL) collected AOR at 2231 h after heavy rains.

THURSTON Co.: H Ave, ca. 0.7 rd km W jct 23rd Rd (42.17699°N, 96.59773°W; WGS 84). 19 May 2017. Samuel M. Tegge and Drew R. Davis. Verified by Travis J. LaDuc. TNHC 103530 (DRD 3973). New county record that fills a gap in the known distribution of this species in Nebraska (Ballinger et al. 2010, *op. cit.*; Fogell 2010, *op. cit.*) and is known from surrounding Burt, Cuming, Dakota, Dixon, and Wayne counties, Nebraska (Ballinger et al. 2010, *op. cit.*; Fogell 2010, *op. cit.*; Davis et al. 2014. *Herpetol. Rev.* 45:99–100; see above) and Monona and Woodbury counties, Iowa (LeClere 2013, *op. cit.*). The nearest known locality for this species is from southeast of Bancroft, Nebraska (TNHC 103528, 103529; see above). Adult female (20.74 g; 56 mm SVL) collected AOR at 2329 h after heavy rains.

Specimens were collected under a Nebraska Game and Parks Commission Scientific and Educational Permit (#1006) issued to DRD.

SAMUEL M. TEGGE, Department of Biology, Western Kentucky University, 1906 College Heights Boulevard, Bowling Green, Kentucky 42101, USA (e-mail: samuel.tegge087@topper.wku.edu); **DREW R. DAVIS**, Department of Biology, University of South Dakota, 414 East Clark Street, Vermillion, South Dakota 57069, USA (e-mail: drew.davis@usd.edu).